




Jet Propulsion Laboratory
California Institute of Technology



Influence of Global Vegetation on Mid-Tropospheric CO₂ Early Results

AIRS Science Team Meeting
April 25, 2012


Thomas S. Pagano, Hai Nguyen, Ed Olsen
California Institute of Technology, Jet Propulsion Laboratory
4800 Oak Grove Drive, Pasadena, CA, USA 91109

tpagano@jpl.nasa.gov, (818) 393-3917, <http://airs.jpl.nasa.gov>

© 2012 California Institute of Technology. Government sponsorship acknowledged.



Introduction




- AIRS Mid-Tropospheric CO₂ shows a high degree of horizontal variability
- Ongoing efforts show AIRS data influenced by global circulation patterns including ENSO and MJO
 - Jiang, X., M. T. Chahine, E. T. Olsen, L. L. Chen, and Y. L. Yung (2010), Interannual variability of mid-tropospheric CO₂ from Atmospheric Infrared Sounder, *Geophys. Res. Lett.*, 37, L13801, doi:10.1029/2010GL042823
 - Li, K. F., B. Tian, D. E. Waliser, Y. L. Yung (2010), Tropical mid-tropospheric CO₂ variability driven by the Madden-Julian oscillation, *PNAS*, 107 (45), 19171-19175, doi:10.1073/pnas.1008222107.
- What is the influence of global vegetation cycle on CO₂ seasonal behavior?
 - Can we correlate mid-trop CO₂ seasonal variability with global vegetation for different regions?
 - For now: First look at zonal averages and Land Vegetation (ocean biomass later)
 - Goal: Sanity Check on AIRS Data Seasonal Cycle, Solicit interest by carbon cycle community
 - Outreach Opportunity: International Workshop on Greenhouse Gas Measurements (IWGGMS) 2012 at CalTech





Jet Propulsion Laboratory
California Institute of Technology

Can Regional Carbon Transport and Storage Be Validated using Satellite Obs?



How well known are
global flux estimates?




$$1 \text{ Pg} = 10^{15} \text{ g} \times (10^{-3} \text{ kg/g}) \times (1\text{T}/907.2 \text{ kg}) = 1.102 \text{ GT}$$



Jet Propulsion Laboratory
California Institute of Technology

Agenda




- CO2 Data
 - AIRS Level 3
 - Creation of a “Climatology”
- Land Vegetation Data: GPP Gross Primary Productivity
 - MODIS Enhanced Vegetation Index
 - MODIS Land Surface Temperature
 - GPP Climatology
- Early Zonal Correlation Results
- Future Work




Jet Propulsion Laboratory
California Institute of Technology



AIRS V5 Mid-Tropospheric CO₂ Shows Horizontal Variability



AIRS Daily CO₂ Yield
1°x1° Spatial Resolution



AIRS Monthly CO₂ Yield
1°x1° Spatial Resolution




Day/Night, Pole-to-Pole, Land/Ocean/Ice, Cloudy/Clear

AIRS CO₂ Data Products Released (2002 to present)
http://airs.jpl.nasa.gov/AIRS_CO2_Data





Jet Propulsion Laboratory
California Institute of Technology

AIRS Mid-Tropospheric CO₂ Representations (“Climatologies”)



Average of L3 Monthly Data by Month over 8 years




Pagano, T. S., Olsen, E. T., Chahine, M. T., Ruzmaikin, A., Nguyen, H., Jiang, X., “[Monthly representations of mid-tropospheric carbon dioxide from the Atmospheric Infrared Sounder](#),” Proc. SPIE 8158-11, San Diego, CA (2011).




Jet Propulsion Laboratory
California Institute of Technology



“Depleted Band of CO₂ Not Due to Surface Vegetation”



Annual Representation of co2midtrop





Downward Flow
of Walker
Circulation





Jet Propulsion Laboratory
California Institute of Technology

Gross Primary Productivity using MODIS Temperature and Greenness (TG)



$$GPP = (\text{scaledEVI} \times \text{scaledLST}) \times m$$

MODIS Night LST

$$\text{scaledLST} = \min \left[\left(\frac{\text{LST}}{30} \right); (2.5 - (0.05 \times \text{LST})) \right]$$


$$\text{scaledEVI} = \text{EVI} - 0.1$$

$$\text{EVI} = G \frac{\rho_{\text{NIR}} - \rho_{\text{Red}}}{\rho_{\text{NIR}} + C_1 \rho_{\text{Red}} - C_2 \rho_{\text{Blue}} + L}$$

$$m = 2.49 - 0.074 \times \text{LST}_{\text{an}} \quad \text{for deciduous sites}$$

$$m = 2.10 - 0.0625 \times \text{LST}_{\text{an}} \quad \text{for evergreen sites.}$$

Using Average m gives < 18% error




Sims, D. et al., A new model of gross primary productivity for North American ecosystems based solely on the enhanced vegetation index and land surface temperature from MODIS, *Remote Sensing of Environment* 112 (2008) 1633–1646



Jet Propulsion Laboratory
California Institute of Technology


Data Sets



- Mirador.gsfc.nasa.gov
- MYDVI.005 MODIS/Aqua Monthly Vegetation Indices Global 1x1 degree
- MYD11CM1N.005 MODIS/Aqua Monthly mean Night-Time Land Surface Temperature at 1x1 degree
- AIRX3C2M.005 AIRS/Aqua Level 3 Monthly CO₂ in the free troposphere (AIRS+AMSU)



MODIS EVI needed to make GPP




- EVI
 - Simple mean EVI (Enhanced Vegetation Index) was calculated from MOD13C2 sds2 CMG 0.05 Deg Monthly EVI only for cells within valid EVI range. Cells with Fill_Values were not included in the analysis. The dataset is produced with full global coverage. Land/water mask is accepted from the original dataset resolution 0.05 degrees. Grid cells with no land surface are assigned the “_FillValue” – 1.0.
- $\text{perc_fill_value} = (\text{Fill_Cell_Count} * 100) / 400$
 - where Fill_Cell_Count is the number of the 0.05 degree cells with the Fill Value within the 1 degree cell, and 400 is the total number of 0.05 degree cells within the aggregated 1 degree cell. The output values are rounded to the nearest integer. Percent Fill Values were calculated for each 1 degree cell. No “_FillValues” are assigned to this layer.
- $\text{perc_good_pixels} = (\text{Count_GOOD_pixel} * 100) / \text{Count_NonFill_Cell}$
 - where Count_GOOD_pixel is the number of 0.05 degree cells flagged as “GOOD” (value 0) quality within the aggregated 1degree cell, Count_NonFill_Cell is the number of 0.05 degree cells within the valid range of pixel reliability values within the aggregated 1 degree cell. The output values are rounded to the nearest integer. Percent GOOD Quality Data are calculated for each 1 degree cell. No “_FillValues” are assigned to this layer.
 - Percent GOOD Quality Data is calculated from MOD13C2 sds13 CMG 0.05 Deg Monthly pixel reliability only for cells within the valid range of values.





Jet Propulsion Laboratory
California Institute of Technology

MODIS EVI Gridded 1x1° Monthly




MODIS EVI for: 6/2003






Jet Propulsion Laboratory
California Institute of Technology



MODIS EVI QC Example



0.05 degree Climate Modeling
Grid global coverage product



Mostly >90% Fill
Mostly >0 QC




$$\text{perc_fill_value} = 100/400 \times 100 = 25\%$$

$$\text{perc_good_pixels} = 70/100 \times 100 = 70\%$$

Ignore perc_fill_value
(for now)



MODIS Night LST needed to make GPP




- **night_lst**

- The dataset contains global monthly-mean day-time and night-time land surface temperature averaged within 1 by 1 degree grid cells. The source for the data is MODIS MOD11C3 product (MODIS Monthly mean land surface temperature at 0.05 degree spatial resolution). The dataset covers the time period strating January 2000.
- The MODIS/Aqua V4 LST/E 8-Day L3 Global CMG product (Short name: MYD11C3) is a monthly composited average, derived from the MYD11C1 daily global product, and stored as clear-sky LST values during a month's period in a 0.05° (5600 meters) geographic CMG. MYD11C3, therefore, inherits all the structural features of its MYD11C1 parent except for the temporal configuration. Please refer to the MYD11C1 product documentation for all algorithm-related details.
- The V4 MYD11C3 product comprises the following Science Data Set (SDS) layers for daytime and nighttime observations: LSTs, quality control assessments, observation times, view zenith angles, clear sky coverages, and emissivities for bands 20, 22, 23, 29, 31, and 32.
- The V4 Aqua/MODIS LST/E products, including MYD11C3, are validated to Stage-1 with well-defined uncertainties over a range of representative conditions. Further details regarding MODIS land product validation for the LST/E products is available from the following URL:
<http://landval.gsfc.nasa.gov/ProductStatus.php?ProductID=MYD11>





Jet Propulsion Laboratory
California Institute of Technology

MODIS Night Land Surface Temperature 1x1°




MODIS Night LST for: 5/2003







Jet Propulsion Laboratory
California Institute of Technology

GPP Quality Control



- Start with Monthly 1x1° EVI and LST Climatologies
- Use QC provided




$$QC_1 = QC_{evi} \times QC_{lst} \times QC_m$$
$$QC_{GPP} = QC_1 \times (GPP \geq 0) \times (LST \geq 0)$$




Jet Propulsion Laboratory
California Institute of Technology


MODIS GPP Climatology Developed using MODIS EVI and LST



MODIS Level 3 GPP Average over 2002-2011




UMT GPP






Jet Propulsion Laboratory
California Institute of Technology

Correlate AIRS CO₂ and MODIS Derived GPP to Qualitatively Assess Influence



$$CO_2 = \overline{CO_2} + \frac{dCO_2}{dt}t + Ao_c \sin(\omega t + \varphi_c)$$







$$GPP = \overline{GPP} + \frac{dGPP}{dt}t + Ao_g \sin(\omega t + \varphi_g)$$

We will compare average global and regional averages, seasonal amplitude and phase
Trend contains high uncertainties that have not yet been quantified (save for later)



Jet Propulsion Laboratory
California Institute of Technology


First look at Averages






Jet Propulsion Laboratory
California Institute of Technology

Seasonal Response Shows Key Differences




- Based on “Climatology” of 2003-2010
- Peak of GPP Response about 3mo after CO₂
- Polar response of CO₂ irregular

GPP




CO₂







Jet Propulsion Laboratory
California Institute of Technology

Seasonal Amplitude Grows as we go Poleward in NH





Using Sine Fit





Jet Propulsion Laboratory
California Institute of Technology


Early Phase Analysis shows Correlation between CO₂ and EVI + Zonal Dependence






Jet Propulsion Laboratory
California Institute of Technology

Zonal Averages of Seasonal Response for GPP and CO₂




Using Max-Min Fit

GPP





CO₂





Jet Propulsion Laboratory
California Institute of Technology



CO₂ and GPP Seasonal Amplitudes Correlated by Zone






Jet Propulsion Laboratory
California Institute of Technology

CO2 and EVI Animation (L. Perkins GSFC/SVS)





Summary, Conclusion, and Future Work



- AIRS Mid-Trop CO₂ has horizontal and seasonal variability that is not well understood at this time
- AIRS Mid-Trop CO₂ Climatology and MODIS GPP Climatology Developed
- Preliminary results show correlation in amplitude of seasonal variability of GPP
 - AIRS CO₂ influenced by seasonal cycle of vegetation
 - Photosynthesis (GPP) influenced by abundance of CO₂
- Future work
 - Extract Principal Component Amplitudes for Improved Correlation Analysis
 - Error Estimates including Spatial Covariance Matrix
 - Add Ocean GPP
 - Examine Regional Correlation
 - Discuss with Carbon Cycle Scientists: IWGGMS 2012